

A TRICHOTOMY FOR A CLASS OF EQUIVALENCE RELATIONS

LONGYUN DING

ABSTRACT. Let $X_n, n \in \mathbb{N}$ be a sequence of non-empty sets, $\psi_n : X_n^2 \rightarrow \mathbb{R}^+$. We consider the relation $E((X_n, \psi_n)_{n \in \mathbb{N}})$ on $\prod_{n \in \mathbb{N}} X_n$ by $(x, y) \in E((X_n, \psi_n)_{n \in \mathbb{N}}) \Leftrightarrow \sum_{n \in \mathbb{N}} \psi_n(x(n), y(n)) < +\infty$. If $E((X_n, \psi_n)_{n \in \mathbb{N}})$ is a Borel equivalence relation, we show a trichotomy that either $\mathbb{R}^{\mathbb{N}}/\ell_1 \leq_B E$, $E_1 \leq_B E$, or $E \leq_B E_0$.

We also prove that, for a rather general case, $E((X_n, \psi_n)_{n \in \mathbb{N}})$ is an equivalence relation iff it is an ℓ_p -like equivalence relation.

1. INTRODUCTION

A topological space is called a *Polish space* if it is separable and completely metrizable. Let X, Y be Polish spaces and E, F equivalence relations on X, Y respectively. A *Borel reduction* of E to F is a Borel function $\theta : X \rightarrow Y$ such that $(x, y) \in E$ iff $(\theta(x), \theta(y)) \in F$, for all $x, y \in X$. We say that E is *Borel reducible* to F , denoted $E \leq_B F$, if there is a Borel reduction of E to F . If $E \leq_B F$ and $F \leq_B E$, we say that E and F are *Borel bireducible* and denote $E \sim_B F$. We refer to [4] and [8] for background on Borel reducibility.

There are several famous dichotomy theorems on Borel reducibility. The first one is the Silver's dichotomy theorem [13].

Theorem 1.1 (Silver). *Let E be a Π_1^1 equivalence relation. Then E has either at most countably many or perfectly many equivalence classes, i.e. $E \leq_B \text{id}(\mathbb{N})$ or $\text{id}(\mathbb{R}) \leq_B E$.*

There are three dichotomy theorems concerning E_0 . Before introducing these theorems, we recall definitions of equivalence relations E_0, E_1, E_0^ω .

- (a) For $x, y \in 2^{\mathbb{N}}$, $(x, y) \in E_0 \Leftrightarrow \exists m \forall n \geq m (x(n) = y(n))$.
- (b) For $x, y \in 2^{\mathbb{N} \times \mathbb{N}}$, $(x, y) \in E_1 \Leftrightarrow \exists m \forall n \geq m \forall k (x(n, k) = y(n, k))$.
- (c) For $x, y \in 2^{\mathbb{N} \times \mathbb{N}}$, $(x, y) \in E_0^\omega \Leftrightarrow \forall k \exists m \forall n \geq m (x(n, k) = y(n, k))$.

Theorem 1.2. *Let E be a Borel equivalence relation. Then*

- (a) (Harrington-Kechris-Louveau [5]) *either $E \leq_B \text{id}(\mathbb{R})$ or $E_0 \leq_B E$;*
- (b) (Kechris-Louveau [9]) *if $E \leq_B E_1$, then $E \leq_B E_0$ or $E \sim_B E_1$;*
- (c) (Hjorth-Kechris [7]) *if $E \leq_B E_0^\omega$, then $E \leq_B E_0$ or $E \sim_B E_0^\omega$.*

Date: January 6, 2010.

2000 Mathematics Subject Classification. Primary 03E15, 54E35, 46A45.

Research partially supported by the National Natural Science Foundation of China (Grant No. 10701044).

Another class of interesting Borel equivalence relations come from classical Banach sequence spaces. Let $p \geq 1$. For $x, y \in \mathbb{R}^{\mathbb{N}}$, $(x, y) \in \mathbb{R}^{\mathbb{N}}/\ell_p \Leftrightarrow x - y \in \ell_p$. It was shown by G. Hjorth [6] that every Borel equivalence relation $E \leq_B \mathbb{R}^{\mathbb{N}}/\ell_1$ is either essentially countable or satisfies $E \sim_B \mathbb{R}^{\mathbb{N}}/\ell_1$. Kanovei asked whether the position of $\mathbb{R}^{\mathbb{N}}/\ell_1$ in the \leq_B -structure is similar with E_1 and E_0^ω (see [8], Question 5.7.5).

Question 1.3 (Kanovei). Does every Borel equivalence relation $E \leq \mathbb{R}^{\mathbb{N}}/\ell_1$ satisfy either $E \leq_B E_0$ or $E \sim_B \mathbb{R}^{\mathbb{N}}/\ell_1$?

Two kinds of ℓ_p -like equivalence relations were introduced by T. Mátrai [11] and the author [1]. (1) Let $f : [0, 1] \rightarrow \mathbb{R}^+$. For $x, y \in [0, 1]^{\mathbb{N}}$, $(x, y) \in E_f \Leftrightarrow \sum_{n \in \mathbb{N}} f(|x(n) - y(n)|) < +\infty$. (2) Let (X_n, d_n) , $n \in \mathbb{N}$ be a sequence of metric spaces, $p \geq 1$. For $(x, y) \in \prod_{n \in \mathbb{N}} X_n$, $(x, y) \in E((X_n, d_n)_{n \in \mathbb{N}}; p) \Leftrightarrow \sum_{n \in \mathbb{N}} d_n(x(n), y(n))^p < +\infty$.

In this paper, we introduce a notion surpassing both (1) and (2). Let X_n , $n \in \mathbb{N}$ be a sequence of non-empty sets, $\psi_n : X_n^2 \rightarrow \mathbb{R}^+$. For $x, y \in \prod_{n \in \mathbb{N}} X_n$, $(x, y) \in E((X_n, \psi_n)_{n \in \mathbb{N}}) \Leftrightarrow \sum_{n \in \mathbb{N}} \psi_n(x(n), y(n)) < +\infty$. Though we did not find a natural necessary and sufficient condition that $E((X_n, \psi_n)_{n \in \mathbb{N}})$ be an equivalence relation, we establish the following trichotomy.

Theorem 1.4. *If $E = E((X_n, \psi_n)_{n \in \mathbb{N}})$ is a Borel equivalence relation, then either $\mathbb{R}^{\mathbb{N}}/\ell_1 \leq_B E$, $E_1 \leq_B E$, or $E \leq_B E_0$.*

From this trichotomy, we can see that Kanovei's problem is valid within equivalence relations of the form $E((X_n, \psi_n)_{n \in \mathbb{N}})$.

It was shown by R. Dougherty and G. Hjorth [2] that, for $p, q \geq 1$, $\mathbb{R}^{\mathbb{N}}/\ell_p \leq_B \mathbb{R}^{\mathbb{N}}/\ell_q$ iff $p \leq q$. We complete a different picture for $0 < p \leq 1$ by showing that $\mathbb{R}^{\mathbb{N}}/\ell_p \sim_B \mathbb{R}^{\mathbb{N}}/\ell_1$.

Via a process of metrization, we prove that, for a rather general case, equivalence relations $E((X_n, \psi_n)_{n \in \mathbb{N}})$ coincide with ℓ_p -like equivalence relations $E((X_n, d_n)_{n \in \mathbb{N}}; p)$.

2. A TRICHOTOMY FOR SUM-LIKE EQUIVALENCE RELATIONS

We denote the set of all non-negative real numbers by \mathbb{R}^+ .

Definition 2.1. Let X_n , $n \in \mathbb{N}$ be a sequence of non-empty sets, $\psi_n : X_n^2 \rightarrow \mathbb{R}^+$. We define a relation $E((X_n, \psi_n)_{n \in \mathbb{N}})$ on $\prod_{n \in \mathbb{N}} X_n$ by

$$(x, y) \in E((X_n, \psi_n)_{n \in \mathbb{N}}) \iff \sum_{n \in \mathbb{N}} \psi_n(x(n), y(n)) < +\infty$$

for $x, y \in \prod_{n \in \mathbb{N}} X_n$. If $(X_n, \psi_n) = (X, \psi)$ for every $n \in \mathbb{N}$, we write $E(X, \psi) = E((X_n, \psi_n)_{n \in \mathbb{N}})$ for the sake of brevity.

It is hard to find a natural necessary and sufficient condition to determine when $E((X_n, \psi_n)_{n \in \mathbb{N}})$ is an equivalence relation. We need the following definition:

Definition 2.2. If $E((X_n, \psi_n)_{n \in \mathbb{N}})$ is an equivalence relation, we call it a *sum-like equivalence relation*. Furthermore, if $X_n, n \in \mathbb{N}$ is a sequence of Polish spaces and every ψ_n is Borel function, it is called a *sum-like Borel equivalence relation*.

The following easy lemma is very useful for the study of sum-like equivalence relations.

Lemma 2.3. *Let $E = E((X_n, \psi_n)_{n \in \mathbb{N}})$ be a sum-like equivalence relation. For $x, y \in \prod_{n \in \mathbb{N}} X_n$, we have*

$$(x, y) \in E \iff \sum_{x(n) \neq y(n)} \psi_n(x(n), y(n)) < +\infty.$$

Proof. Since E is an equivalence relation, $(x, x) \in E$. It follows that

$$\sum_{x(n)=y(n)} \psi_n(x(n), y(n)) \leq \sum_{n \in \mathbb{N}} \psi_n(x(n), x(n)) < +\infty.$$

Then the lemma follows. \square

Definition 2.4. Let $(X_n, d_n), n \in \mathbb{N}$ be a sequence of pseudo-metric spaces. For $p \geq 1$, an ℓ_p -like equivalence relation $E((X_n, d_n)_{n \in \mathbb{N}}; p)$ is $E((X_n, \psi_n)_{n \in \mathbb{N}})$ where $\psi_n(u, v) = d_n(u, v)^p$ for $u, v \in X_n$. If $(X_n, d_n) = (X, d)$ for every $n \in \mathbb{N}$, we write $E(X, d; p) = E((X_n, d_n)_{n \in \mathbb{N}}; p)$ for the sake of brevity.

Proposition 2.5. *Let $E((X_n, d_n)_{n \in \mathbb{N}}; p)$ be an ℓ_p -like equivalence relation. Then there exists metric d'_n on X_n for each n such that $E((X_n, d'_n)_{n \in \mathbb{N}}; p) = E((X_n, d_n)_{n \in \mathbb{N}}; p)$.*

Proof. We can see that following d'_n 's meet the requirements.

$$d'_n(u, v) = \begin{cases} 0, & u = v, \\ 2^{-n}, & u \neq v, d_n(u, v) \leq 2^{-n}, \\ d_n(u, v), & d_n(u, v) > 2^{-n}, \end{cases}$$

for $u, v \in X_n$. \square

Proposition 2.6. *Let $E_i, i \in \mathbb{N}$ be a sequence of Borel equivalence relations. Then for every $p \geq 1$ there is an ℓ_p -like Borel equivalence relation $E = E((X_n, d_n)_{n \in \mathbb{N}}; p)$ such that $E_i \leq_B E$ for $n \in \mathbb{N}$.*

Proof. For $i \in \mathbb{N}$, let E_i be a Borel equivalence relation on a Polish space Y_i . We define $\chi_i : Y_i^2 \rightarrow \mathbb{R}^+$ by

$$\chi_i(u, v) = \begin{cases} 0, & (u, v) \in E_i, \\ 1, & (u, v) \notin E_i. \end{cases}$$

Now fix a bijection $\langle \cdot, \cdot \rangle : \mathbb{N}^2 \rightarrow \mathbb{N}$. For every $i, j \in \mathbb{N}$, if $n = \langle i, j \rangle$, we denote $X_n = Y_i$ and $d_n = \chi_i$. It is easy to verify that $E = E((X_n, d_n)_{n \in \mathbb{N}}; p)$ is an ℓ_p -like Borel equivalence relation. For $i \in \mathbb{N}$, define $\theta_i : Y_i \rightarrow \prod_{n \in \mathbb{N}} X_n$ by

$$\theta_i(u)(\langle k, j \rangle) = \begin{cases} u, & k = i, \\ a_k, & k \neq i, \end{cases}$$

where $a_k \in Y_k$ is independent of u . Clearly θ_i is a Borel reduction of E_i to E . \square

Now suppose that $E = E((X_n, \psi_n)_{n \in \mathbb{N}})$ is a sum-like Borel equivalence relation. Its position in the \leq_B -structure of Borel equivalence relations is determined by the following condition:

($\ell 1$) $\forall c > 0 \exists x, y \in \prod_{n \in \mathbb{N}} X_n$ such that $\exists m \forall n > m (\psi_n(x(n), y(n)) < c)$ and

$$\sum_{n \in \mathbb{N}} \psi_n(x(n), y(n)) = +\infty.$$

Lemma 2.7. *If ($\ell 1$) holds, then $\mathbb{R}^{\mathbb{N}}/\ell_1 \leq_B E$.*

Proof. Firstly, we show a well known fact: $\mathbb{R}^{\mathbb{N}}/\ell_1 \leq_B [0, 1]^{\mathbb{N}}/\ell_1$. Fix a bijection $\langle \cdot, \cdot \rangle' : \mathbb{N} \times \mathbb{Z} \rightarrow \mathbb{N}$. For $z \in \mathbb{R}^{\mathbb{N}}$, we define $\theta'(z) \in [0, 1]^{\mathbb{N}}$ as

$$\theta'(z)(\langle m, k \rangle') = \begin{cases} 0, & z(m) < k, \\ z(m) - k, & k \leq z(m) < k + 1, \\ 1, & k + 1 \leq z(m). \end{cases}$$

Then θ' witness that $\mathbb{R}^{\mathbb{N}}/\ell_1 \leq_B [0, 1]^{\mathbb{N}}/\ell_1$.

Secondly, we construct a reduction of $[0, 1]^{\mathbb{N}}/\ell_1$ to E .

For $l \in \mathbb{N}$, by condition ($\ell 1$), there exist $x_l, y_l \in \prod_{n \in \mathbb{N}} X_n$ such that $\exists m \forall n > m (\psi_n(x_l(n), y_l(n)) < 2^{-l})$ and $\sum_{n \in \mathbb{N}} \psi_n(x_l(n), y_l(n)) = +\infty$. Then we can select two sequences of natural numbers $(i_l)_{l \in \mathbb{N}}, (j_l)_{l \in \mathbb{N}}$ satisfying that

- (i) $i_l < j_l < i_{l+1}$ for $l \in \mathbb{N}$;
- (ii) $\psi_n(x_l(n), y_l(n)) < 2^{-l}$ for $i_l \leq n \leq j_l$;
- (iii) $1 \leq \sum_{i_l \leq n \leq j_l} \psi_n(x_l(n), y_l(n)) < 1 + 2^{-l}$.

Fix an element $a_n \in X_n$ for every $n \in \mathbb{N}$. For $z \in [0, 1]^{\mathbb{N}}$, we define $\vartheta(z) \in \prod_{n \in \mathbb{N}} X_n$ by

$$\vartheta(z)(n) = \begin{cases} x_l(n), & i_l \leq n \leq j_l, \sum_{i_l \leq m \leq n} \psi_m(x_l(m), y_l(m)) \leq z(l); \\ y_l(n), & i_l \leq n \leq j_l, \sum_{i_l \leq m \leq n} \psi_m(x_l(m), y_l(m)) > z(l); \\ a_n, & \text{otherwise.} \end{cases}$$

Note that, for $z, w \in [0, 1]^{\mathbb{N}}$ and $l \in \mathbb{N}$, we have

$$|z(l) - w(l)| - 2^{-l} < \sum_{\substack{i_l \leq n \leq j_l \\ \vartheta(z)(n) \neq \vartheta(w)(n)}} \psi_n(\vartheta(z)(n), \vartheta(w)(n)) < |z(l) - w(l)| + 2^{-l}.$$

Therefore, by Lemma 2.3,

$$\begin{aligned} (\vartheta(z), \vartheta(w)) \in E & \iff \sum_{\vartheta(z)(n) \neq \vartheta(w)(n)} \psi_n(\vartheta(z)(n), \vartheta(w)(n)) < +\infty \\ & \iff \sum_{l \in \mathbb{N}} |z(l) - w(l)| < +\infty \\ & \iff z - w \in \ell_1. \end{aligned}$$

It follows that $[0, 1]^{\mathbb{N}}/\ell_1 \leq_B E$. \square

If $(\ell 1)$ fails, then there exists $c > 0$ such that

$$\exists m \forall n > m (\psi_n(x(n), y(n)) < c) \Rightarrow \sum_{n \in \mathbb{N}} \psi_n(x(n), y(n)) < +\infty$$

for $x, y \in \prod_{n \in \mathbb{N}} X_n$. Now we denote

$$F_n = \{(u, v) \in X_n^2 : \psi_n(u, v) < c\}.$$

Lemma 2.8. *If $(\ell 1)$ fails, let $c > 0$ and $F_n, n \in \mathbb{N}$ be defined as above. Then*

(1) *for $x, y \in \prod_{n \in \mathbb{N}} X_n$,*

$$(x, y) \in E \iff \exists m \forall n > m ((x(n), y(n)) \in F_n);$$

(2) *there exists N_0 such that, for $n > N_0$, F_n is a Borel equivalence relation on X_n .*

Proof. Since $(\ell 1)$ fails, clause (1) is trivial.

(2) Firstly, assume for contradiction that, there exist a strictly increasing sequence of natural numbers $(n_k)_{k \in \mathbb{N}}$ and $u_k \in X_{n_k}$ such that $\psi_{n_k}(u_k, u_k) \geq c$. Select an $x \in \prod_{n \in \mathbb{N}} X_n$ with $x(n_k) = u_k$ for $k \in \mathbb{N}$. Then we have $\sum_{n \in \mathbb{N}} \psi_n(x(n), x(n)) = +\infty$. This is impossible, since E is an equivalence relation. So there exists N_1 such that, for $n > N_1, u \in X_n$, we have $\psi_n(u, u) < c$, i.e. $(u, u) \in F_n$.

Secondly, assume for contradiction that, there exist a strictly increasing sequence of natural numbers $(n_k)_{k \in \mathbb{N}}$ and $u_k, v_k \in X_{n_k}$ such that $\psi_{n_k}(u_k, v_k) < c, \psi_{n_k}(v_k, u_k) \geq c$. Select $x, y \in \prod_{n \in \mathbb{N}} X_n$ such that $x(n_k) = u_k, y(n_k) = v_k$ for $k \in \mathbb{N}$ and $x(n) = y(n)$ for other n . We have $\psi_n(x(n), y(n)) < c$ for $n > N_1$. Since $(\ell 1)$ fails, $\sum_{n \in \mathbb{N}} \psi_n(x(n), y(n)) < +\infty$, i.e. $(x, y) \in E$. Clearly $\sum_{n \in \mathbb{N}} \psi_n(y(n), x(n)) = +\infty$, so $(y, x) \notin E$. A contradiction! Hence there exists N_2 such that, for $n > N_2, u, v \in X_n$, we have $\psi_n(u, v) < c \Rightarrow \psi_n(v, u) < c$, i.e. $(u, v) \in F_n \Rightarrow (v, u) \in F_n$.

With a similar argument, we can prove that there exists N_3 such that, for $n > N_3, u, v, r \in X_n$, we have $(u, v), (v, r) \in F_n \Rightarrow (u, r) \in F_n$.

In summary, for $n > N_0 = \max\{N_1, N_2, N_3\}$, F_n is an equivalence relation. Since ψ_n is Borel, so is F_n . \square

Recall that $E_0(\mathbb{N})$ is an equivalence relation on $\mathbb{N}^{\mathbb{N}}$ similar to E_0 on $2^{\mathbb{N}}$. For $x, y \in \mathbb{N}^{\mathbb{N}}$, $(x, y) \in E_0(\mathbb{N}) \Leftrightarrow \exists m \forall n > m (x(n) = y(n))$. It is well known that $E_0 \sim_B E_0(\mathbb{N})$ (see Proposition 6.1.2 of [4]).

Lemma 2.9. *For any equivalence relation E on $\prod_{n \in \mathbb{N}} X_n$, if there is a sequence $F_n \subseteq X_n^2, n \in \mathbb{N}$ such that (1) and (2) of Lemma 2.8 hold, then either $E_1 \leq_B E$, $E \sim_B E_0$, or E is trivial, i.e. all elements in $\prod_{n \in \mathbb{N}} X_n$ are equivalent.*

Proof. For $n > N_0$, since F_n is a Borel equivalence relation on X_n , from the Silver dichotomy theorem [13], either there are at most countably many F_n -equivalence classes or there are perfectly many F_n -equivalence classes.

Case 1. There exists a strictly increasing sequence of natural numbers $(n_k)_{k \in \mathbb{N}}$ such that there are perfectly many F_{n_k} -equivalence classes. Then there is a continuous embedding $h_k : 2^{\mathbb{N}} \rightarrow X_{n_k}$ for every $k \in \mathbb{N}$ such that $(h_k(z), h_k(w)) \in F_{n_k}$ iff $z = w$. Define $\theta : 2^{\mathbb{N} \times \mathbb{N}} \rightarrow \prod_{n \in \mathbb{N}} X_n$ by

$$\theta(x)(n) = \begin{cases} h_k(x(k, \cdot)), & n = n_k, \\ a_n, & \text{otherwise,} \end{cases}$$

where $a_n \in X_n$ is independent of x . By Lemma 2.8.(1), it is straightforward to check that θ is a reduction of E_1 to E .

Case 2. There exists N such that, for $n > N$, F_n has only one equivalence class. From Lemma 2.8.(1), we see that E is trivial.

Case 3. If case 1 fails, then there exists N' such that, for $n > N'$, F_n has at most countably many equivalence classes. So $E \leq_B E_0(\mathbb{N})$. If case 2 fails, then there exists a strictly increasing sequence of natural numbers $(n_k)_{k \in \mathbb{N}}$ such that F_{n_k} has more than one equivalence class. Thus $E_0 \leq_B E$. Since $E_0 \sim_B E_0(\mathbb{N})$, we have $E \sim_B E_0$. \square

Now we have already completed the proof of the following trichotomy.

Theorem 2.10. *Let $E = E((X_n, \psi_n)_{n \in \mathbb{N}})$ be a sum-like Borel equivalence relation. Then either $\mathbb{R}^{\mathbb{N}}/\ell_1 \leq_B E$, $E_1 \leq_B E$, or $E \leq_B E_0$.*

Corollary 2.11. *Let $E = E((X_n, \psi_n)_{n \in \mathbb{N}})$ be a sum-like Borel equivalence relation. If $E \leq_B \mathbb{R}^{\mathbb{N}}/\ell_1$, then either $E \leq_B E_0$ or $E \sim_B \mathbb{R}^{\mathbb{N}}/\ell_1$.*

Proof. It is well known that $E_1 \not\leq_B \mathbb{R}^{\mathbb{N}}/\ell_1$ (see [9] Theorem 4.2), so if $E \leq_B \mathbb{R}^{\mathbb{N}}/\ell_1$, then $E_1 \not\leq_B E$. Hence the corollary follows. \square

It was shown by R. Dougherty and G. Hjorth [2] that, for $p, q \geq 1$,

$$\mathbb{R}^{\mathbb{N}}/\ell_p \leq_B \mathbb{R}^{\mathbb{N}}/\ell_q \iff p \leq q.$$

The following corollary shows that, for $p \leq 1$, the situation is different.

Corollary 2.12. *For $0 < p \leq 1$, we have $\mathbb{R}^{\mathbb{N}}/\ell_p \sim_B \mathbb{R}^{\mathbb{N}}/\ell_1$.*

Proof. Note that $\mathbb{R}^{\mathbb{N}}/\ell_p = E(\mathbb{R}, \psi)$ where $\psi(u, v) = |u - v|^p$ for $u, v \in \mathbb{R}$. It is easy to see that (ℓ_1) holds for $E(\mathbb{R}, \psi)$, so $\mathbb{R}^{\mathbb{N}}/\ell_1 \leq_B \mathbb{R}^{\mathbb{N}}/\ell_p$.

For the other direction, we claim that $\mathbb{R}^{\mathbb{N}}/\ell_p \leq_B \mathbb{R}^{\mathbb{N}}/\ell_q$ for $0 < p < q \leq 1$.

We sketch the proof for Theorem 1.1 of [2], that $\mathbb{R}^{\mathbb{N}}/\ell_p \leq_B \mathbb{R}^{\mathbb{N}}/\ell_q$ for $1 \leq p < q$, and check that it is also valid for $0 < p < q \leq 1$.

It will suffice to prove for the case $0 < \frac{q}{2} < p < q \leq 1$. We denote $\rho = \frac{p}{q}$ and $r = 4^{-\rho}$. Then $\frac{1}{2} < \rho < 1$ and $\frac{1}{4} < r < \frac{1}{2}$. In [2] p. 1838, the authors constructed a continuous function $\bar{K}_r : \mathbb{R} \rightarrow \mathbb{R}^2$, and proved that there are $m', M' > 0$ such that

$$m'|s - t|^\rho \leq \|\bar{K}_r(s) - \bar{K}_r(t)\|_2 \leq M'|s - t|^\rho,$$

for $s, t \in [i - 1, i + 1], i \in \mathbb{Z}$. And $\|\bar{K}_r(s) - \bar{K}_r(t)\|_2 \geq 1$ if s, t are not in the same interval $[i - 1, i + 1], i \in \mathbb{Z}$.

Define a mapping $\theta : \mathbb{R}^{\mathbb{N}} \rightarrow \mathbb{R}^{\mathbb{N}}$ such that, for $x \in \mathbb{R}^{\mathbb{N}}$ and $k \in \mathbb{N}$,

$$\bar{K}_r(x(k)) = (\theta(x)(2k), \theta(x)(2k+1)).$$

For $w = (s, t) \in \mathbb{R}^2$, denote $\|w\|_q = (|s|^q + |t|^q)^{\frac{1}{q}}$. Note that

$$\frac{1}{\sqrt{2}}\|w\|_2 \leq \|w\|_{\infty} \leq \|w\|_q \leq 2^{\frac{1}{q}}\|w\|_{\infty} \leq 2^{\frac{1}{q}}\|w\|_2.$$

For $x, y \in \mathbb{R}^{\mathbb{N}}$, we have

$$\begin{aligned} \theta(x) - \theta(y) \in \ell_q &\iff \sum_{k \in \mathbb{N}} \|\bar{K}_r(x(k)) - \bar{K}_r(y(k))\|_q^q < +\infty \\ &\iff \sum_{k \in \mathbb{N}} \|\bar{K}_r(x(k)) - \bar{K}_r(y(k))\|_2^q < +\infty \\ &\iff \sum_{k \in \mathbb{N}} |x(k) - y(k)|^p < +\infty \\ &\iff x - y \in \ell_p. \end{aligned}$$

Thus, θ is a reduction of $\mathbb{R}^{\mathbb{N}}/\ell_p$ to $\mathbb{R}^{\mathbb{N}}/\ell_q$. \square

3. METRIZATION

In this section, we show that a sum-like equivalence relation $E((X_n, \psi_n)_{n \in \mathbb{N}})$ coincides with an ℓ_p -like equivalence relation if the following conditions hold.

(m1) Denote $X = \bigcup_{n \in \mathbb{N}} X_n$. There is a unique function $\psi : X^2 \rightarrow \mathbb{R}^+$ such that $\psi_n = \psi \upharpoonright X_n^2$ and $\psi(u, v) = 1$ if no X_n contains both u, v .

(m2) For any $u, v, r \in X$, if $u, v, r \in X_n$, then there exists $m > n$ such that $u, v, r \in X_m$.

Let $\bar{\psi}_n = \min\{\psi_n, 1\}$. We can see that $E((X_n, \bar{\psi}_n)_{n \in \mathbb{N}}) = E((X_n, \psi_n)_{n \in \mathbb{N}})$. Therefore, we may assume $\psi_n \leq 1$ if needed.

Lemma 3.1. *Let $(X_n, \psi_n)_{n \in \mathbb{N}}$ satisfy (m1) and (m2). If $\psi_n \leq 1$, then $E((X_n, \psi_n)_{n \in \mathbb{N}})$ is an equivalence relation iff the following conditions hold:*

- (i) $\psi(u, u) = 0$ for $u \in X$;
- (ii) there is a $C \geq 1$ such that for $u, v, r \in X$,

$$\psi(v, u) \leq C\psi(u, v); \quad \psi(u, r) \leq C(\psi(u, v) + \psi(v, r)).$$

Proof. If conditions (i) and (ii) hold, it is trivial that $E((X_n, \psi_n)_{n \in \mathbb{N}})$ is an equivalence relation. We only need to prove the other direction.

Now assume that $E((X_n, \psi_n)_{n \in \mathbb{N}})$ is an equivalence relation.

Firstly, for any $u \in X$, by (m2), there is an infinite set $I \subseteq \mathbb{N}$ such that $u \in X_n$ for $n \in I$. Let $x \in \prod_{n \in \mathbb{N}} X_n$ with $x(n) = u$ for $n \in I$. Since $(x, x) \in E((X_n, \psi_n)_{n \in \mathbb{N}})$, we have $\psi(u, u) = 0$.

Secondly, for $u, v \in X$, if $\psi(u, v) = 0$, we claim that $\psi(v, u) = 0$. By (m2), there is an infinite set $I \subseteq \mathbb{N}$ such that $u, v \in X_n$ for $n \in I$. Therefore, for any $x, y \in \prod_{n \in \mathbb{N}} X_n$, if $x(n) = u, y(n) = v$ for $n \in I$ and $x(n) = y(n)$ for $n \notin I$, we have $(x, y) \in E((X_n, \psi_n)_{n \in \mathbb{N}})$. Hence $(y, x) \in E((X_n, \psi_n)_{n \in \mathbb{N}})$. It follows that $\sum_{n \in I} \psi(v, u) = \sum_{n \in I} \psi(y(n), x(n)) < +\infty$. Thus $\psi(v, u) = 0$.

Now assume for contradiction that, for every $k \in \mathbb{N}$ there are $u_k, v_k \in X$ such that $\psi(v_k, u_k) > 2^k \psi(u_k, v_k) > 0$. Since $\psi_n \leq 1$, $0 < \psi(u_k, v_k) < 2^{-k}$. By (m2), there are infinitely many n such that $u_k, v_k \in X_n$.

Select a finite set $I_k \subseteq \mathbb{N}$ for every k satisfying that

- (i) $u_k, v_k \in X_n$ for $n \in I_k$;
- (ii) $2^{-k} \leq |I_k| \psi(u_k, v_k) \leq 2^{-(k-1)}$;
- (iii) if $k_1 < k_2$, then $\max I_{k_1} < \min I_{k_2}$.

Now we define $x, y \in \prod_{n \in \mathbb{N}} X_n$ by

$$\begin{cases} x(n) = u_k, y(n) = v_k, & n \in I_k, k \in \mathbb{N}, \\ x(n) = y(n) = a_n, & \text{otherwise,} \end{cases}$$

where $a_n \in X_n$ is independent of x and y . Then we have

$$\sum_{n \in \mathbb{N}} \psi(x(n), y(n)) = \sum_{k \in \mathbb{N}} |I_k| \psi(u_k, v_k) \leq \sum_{k \in \mathbb{N}} 2^{-(k-1)} < +\infty,$$

so $(x, y) \in E((X_n, \psi_n)_{n \in \mathbb{N}})$. On the other hand, we have

$$\sum_{n \in \mathbb{N}} \psi(y(n), x(n)) = \sum_{k \in \mathbb{N}} |I_k| \psi(v_k, u_k) > \sum_{k \in \mathbb{N}} 2^k |I_k| \psi(u_k, v_k) \geq \sum_{k \in \mathbb{N}} 1 = +\infty,$$

so $(y, x) \notin E((X_n, \psi_n)_{n \in \mathbb{N}})$. A Contradiction! We complete the proof of that there is $C_1 \geq 1$ such that $\psi(u, v) \leq C_1 \psi(v, u)$ for $u, v \in X$.

With a similar argument, we can prove that there is $C_2 \geq 1$ such that $\psi(u, r) \leq C_2(\psi(u, v) + \psi(v, r))$ for $u, v, r \in X$. \square

Lemma 3.2. *Let $E((X_n, \psi_n)_{n \in \mathbb{N}}), E((X_n, \varphi_n)_{n \in \mathbb{N}})$ be two sum-like equivalence relations, both satisfying (m1) and (m2). If $\varphi_n \leq 1$, then*

$$\begin{aligned} & E((X_n, \psi_n)_{n \in \mathbb{N}}) \subseteq E((X_n, \varphi_n)_{n \in \mathbb{N}}) \\ \iff & \exists A \geq 1 \forall u, v \in X (\varphi(u, v) \leq A \psi(u, v)). \end{aligned}$$

Proof. “ \Leftarrow ” is trivial. “ \Rightarrow ” follows similarly as the proof of Lemma 3.1. \square

Corollary 3.3. *Let $E((X_n, \psi_n)_{n \in \mathbb{N}}), E((X_n, \varphi_n)_{n \in \mathbb{N}})$ be two sum-like equivalence relations, both satisfying (m1) and (m2). If $\psi_n, \varphi_n \leq 1$, then*

$$\begin{aligned} & E((X_n, \psi_n)_{n \in \mathbb{N}}) = E((X_n, \varphi_n)_{n \in \mathbb{N}}) \\ \iff & \exists A \geq 1 \forall u, v \in X (A^{-1} \psi(u, v) \leq \varphi(u, v) \leq A \psi(u, v)). \end{aligned}$$

Before introducing the Metrization lemma, we recall several basic notions on relations. Let X be a non-empty set, we denote $\Delta(X) = \{(u, u) : u \in X\}$. A subset $U \subseteq X^2$ is called symmetric if $(u, v) \in U \Rightarrow (v, u) \in U$ for $u, v \in X$. For $U, V \subseteq X^2$ we define $U \circ V \subseteq X^2$ by

$$(u, r) \in U \circ V \iff \exists v ((u, v) \in U, (v, r) \in V) \quad (\forall u, v, r \in X).$$

Lemma 3.4 (Metrization lemma [10], p. 185). *Let $U_n, n \in \mathbb{N}$ be a sequence of subsets of X^2 such that*

- (i) $U_0 = X^2$;
- (ii) each U_n is symmetric and $\Delta(X) \subseteq U_n$;
- (iii) $U_{n+1} \circ U_{n+1} \subseteq U_n$ for each n .

Then there is a pseudo-metric d on X satisfying that

$$U_n \subseteq \{(u, v) : d(u, v) < 2^{-n}\} \subseteq U_{n-1} \quad (\forall n \geq 1).$$

Theorem 3.5. *Let $(X_n, \psi_n)_{n \in \mathbb{N}}$ satisfy (m1) and (m2). Then $E((X_n, \psi_n)_{n \in \mathbb{N}})$ is an equivalence relation iff it is an ℓ_p -like equivalence relation.*

Proof. Assume that $E((X_n, \psi_n)_{n \in \mathbb{N}})$ is an equivalence relation. Without loss of generality, we may assume that $\psi_n \leq 1$. From Lemma 3.1, there is $C \geq 1$ such that, for $u, v, r \in X$,

$$\psi(v, u) \leq C\psi(u, v) \text{ and } \psi(u, r) \leq C(\psi(u, v) + \psi(v, r)).$$

Therefore, if $\psi(u, v), \psi(v, r), \psi(r, s) < \varepsilon$, then

$$\psi(u, s) \leq C(\psi(u, v) + C(\psi(v, r) + \psi(r, s))) < (2C^2 + C)\varepsilon.$$

Now denote $B = 2C^2 + C$. We define $U_0 = X^2$ and

$$U_n = \{(u, v) : \psi(u, v) < B^{-n}, \psi(v, u) < B^{-n}\} \quad (n \geq 1).$$

It follows that, for each n , U_n is symmetric and $U_{n+1} \circ U_{n+1} \circ U_{n+1} \subseteq U_n$. By Lemma 3.1.(i), $\Delta(X) \subseteq U_n$. Then the metrization lemma gives a pseudo-metric d on X such that $U_n \subseteq \{(u, v) : d(u, v) < 2^{-n}\} \subseteq U_{n-1}$.

It is easy to check that, $d(u, v) = 0$ iff $\psi(u, v) = 0$. If $\psi(u, v) \geq B^{-2}$, then $(u, v) \notin U_2$, $d(u, v) \geq 2^{-3}$.

Denote $p = \log_2 B \geq 1$. If $0 < \psi(u, v) < B^{-2}$, assume that $B^{-(n+1)} \leq \psi(u, v) < B^{-n}$ for some $n \geq 2$. Then $\psi(v, u) < CB^{-n} < B^{-(n-1)}$. It follows that $(u, v) \in U_{n-1}$, $(u, v) \notin U_{n+1}$, $2^{-(n+2)} \leq d(u, v) < 2^{-(n-1)}$. So

$$B^{-2}d(u, v)^p < B^{-2}(2^{-(n-1)})^p \leq \psi(u, v) < B^2(2^{-(n+2)})^p \leq B^2d(u, v)^p.$$

Therefore, we have $E((X_n, \psi_n)_{n \in \mathbb{N}}) = E((X_n, d \upharpoonright X_n); p)$. \square

Corollary 3.6. *Let $(X_n, \psi_n)_{n \in \mathbb{N}}$ satisfy that, for $n < m$, $X_n \subseteq X_m$ and $\psi_n = \psi_m \upharpoonright X_n^2$. Then $E((X_n, \psi_n)_{n \in \mathbb{N}})$ is an equivalence relation iff it is an ℓ_p -like equivalence relation.*

In particular, $E(X, \psi)$ is an equivalence relation iff there are pseudo-metric d on X and $p \geq 1$ such that $E(X, \psi) = E(X, d; p)$.

4. FURTHER REMARKS

Let us consider a special case of sum-like equivalence relation. Let $\psi_f(u, v) = f(|u - v|)$ where $f : \mathbb{R}^+ \rightarrow \mathbb{R}^+$. Then conditions (i) and (ii) for $E(\mathbb{R}, \psi_f)$ in Lemma 3.1 read as (see also [11], Proposition 2).

- (i) $f(0) = 0$;
- (ii) there is a $C \geq 1$ such that for $s, t \in \mathbb{R}^+$,

$$f(s + t) \leq C(f(s) + f(t)), \quad f(s) \leq C(f(s + t) + f(t)).$$

Denote $\mathcal{N}_f = \{x \in \mathbb{R}^{\mathbb{N}} : \sum_{n \in \mathbb{N}} f(|x(n)|) < +\infty\}$. Then $E(\mathbb{R}, \psi_f)$ is an equivalence relation iff \mathcal{N}_f is a subgroup of $(\mathbb{R}^{\mathbb{N}}, +)$. Furthermore, another interesting problem is to determine when \mathcal{N}_f is a linear subspace of $\mathbb{R}^{\mathbb{N}}$. This problem was studied by S. Mazur and W. Orlicz (see [12], 1.7). It was also considered in [12] that when \mathcal{N}_f 's are Banach spaces.

Theorem 4.1 (Mazur-Orlicz). *Let $f : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ satisfy that, as $n \rightarrow \infty$, $f(t_n) \rightarrow 0$ iff $t_n \rightarrow 0$. The necessary and sufficient condition for \mathcal{N}_f to be a linear space is that*

- (a) *there exist constants $C > 0, \varepsilon > 0$ such that $f(s+t) \leq C(f(s) + f(t))$ for $s, t < \varepsilon$;*
- (b) *for every $\rho > 0$ there are constants $D > 0, \delta > 0$ such that $f(s) \leq Df(t)$ for $t < \delta, s < \rho t$.*

Note that for $t_n \geq \min\{\varepsilon, \delta\} > 0$, $n \in \mathbb{N}$, we have $f(t_n) \not\rightarrow 0$. Thus there is $c > 0$ such that $f(t) \geq c$ for $t \geq \min\{\varepsilon, \delta\}$. If we assume that $f \leq 1$, then conditions (a) and (b) in this theorem turn to

- (a)' *there exists a constant $C' > 0$ such that $f(2s) \leq C'f(s)$ for $s \in \mathbb{R}^+$;*
- (b)' *there exists a constant $D' > 0$ such that $f(s) \leq D'f(t)$ for $s < t$.*

For almost all known examples of sum-like equivalence relations $E(\mathbb{R}, \psi_f)$, \mathcal{N}_f 's are linear spaces. In the end, we present an example in which \mathcal{N}_f is not linear as follows.

Example 4.2. Let $g : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ be an increasing function with $g(0) = 0$, such that $g'(t)$ is decreasing with $\lim_{t \rightarrow 0} g'(t) = +\infty$. For example, $g(x) = \sqrt{x}$ is such a function. Let $(a_n)_{n \in \mathbb{N}}$ be a strictly decreasing sequence of positive numbers with $\lim_{n \rightarrow \infty} a_n = 0$.

Denote $k_n = \frac{g(a_n)}{a_n}$. Then $k_n < k_{n+1}$. We consider equations $y = k_n x$ and $y - g(a_{n+1}) = -k_{n+1}(x - a_{n+1})$. Their solution is $(b_n, k_n b_n)$ where $b_n = \frac{2k_{n+1}a_{n+1}}{k_n + k_{n+1}}$. We can see that $a_{n+1} < b_n < a_n$.

Now define $f : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ by

$$f(t) = \begin{cases} 0, & t = 0, \\ -k_{n+1}(t - a_{n+1}) + g(a_{n+1}), & a_{n+1} \leq t < b_n, \\ k_n t, & b_n \leq t < a_n, \\ g(a_0), & a_0 \leq t. \end{cases}$$

It is easy to check that, for $s, t \in \mathbb{R}^+$,

$$f(s+t) \leq f(s) + f(t), \quad f(s) \leq f(s+t) + f(t).$$

Note that $\frac{f(a_{n+1})}{f(b_n)} = \frac{1}{2} \left(1 + \frac{k_{n+1}}{k_n} \right)$. Since $\lim_{t \rightarrow 0} \frac{g(t)}{t} = \lim_{t \rightarrow 0} g'(t) = +\infty$, we can find a sequence $(a_n)_{n \in \mathbb{N}}$ such that $\frac{k_{n+1}}{k_n} \rightarrow +\infty$, $\frac{f(a_{n+1})}{f(b_n)} \rightarrow +\infty$ as $n \rightarrow \infty$. Then condition (b) in Theorem 4.1 fails.

REFERENCES

- [1] L. Ding, *Borel reducibility and Hölder(α) embeddability between Banach spaces*, preprint, available at <http://arxiv.org/abs/0912.1912>.
- [2] R. Dougherty and G. Hjorth, *Reducibility and nonreducibility between ℓ^p equivalence relations*, Trans. Amer. Math. Soc. 351 (1999) 1835–1844.
- [3] I. Farah, *Basis problem for turbulent actions II: c_0 -equalities*, Proc. London Math. Soc. 82 (2001), no. 3, 1–30.
- [4] S. Gao, *Invariant Descriptive Set Theory*, Monographs and Textbooks in Pure and Applied Mathematics, vol. 293, CRC Press, 2008.

- [5] L. A. Harrington, A. S. Kechris, and A. Louveau, *A Glimm-Effros dichotomy for Borel equivalence relations*, J. Amer. Math. Soc. 3 (1990), no. 4, 903–928.
- [6] G. Hjorth, *Actions by the classical Banach spaces*, J. Symbolic Logic 65 (2000), no. 1, 392–420.
- [7] G. Hjorth and A. S. Kechris, *New dichotomies for Borel equivalence relations*, Bull. Symbolic Logic 3 (1997), no. 3, 329–346.
- [8] V. Kanovei, *Borel Equivalence Relations: Structure and Classification*, University Lecture Series, vol. 44, A. M. S., 2008.
- [9] A. S. Kechris and A. Louveau, *The classification of hypersmooth Borel equivalence relations*, J. Amer. Math. Soc. 10 (1997), no. 1, 215–242.
- [10] J. L. Kelley, *General Topology*, D. Van Nostrand Company, Inc., 1955.
- [11] T. Mátrai, *On ℓ_p -like equivalence relations*, Real Anal. Exchange 34 (2008/09), no. 2, 377–412.
- [12] S. Mazur and W. Orlicz, *On some classes of linear spaces*, Studia Math., 17 (1958), 97–119.
- [13] J. H. Silver, *Counting the number of equivalence classes of Borel and coanalytic equivalence relations*, Ann. Math. Logic 18 (1980), no. 1, 1–28.

SCHOOL OF MATHEMATICAL SCIENCES AND LPMC, NANKAI UNIVERSITY, TIANJIN, 300071, P.R.CHINA

E-mail address: dinglongyun@gmail.com